



CLIFFORD M. DAVIDSON  
LESLIE B. DAVIDSON  
CARY S. KAPPEL  
WILLIAM C. GEHRIS  
MOREY B. WILDES  
ROBERT J. PARADISO  
ERIK R. SWANSON\*\*  
THOMAS P. CANTY\*\*



SEP 29 2004



FELIX L. D'ARIENZO, JR.  
STEPHANIE HSIEH

DAVID G. KNASIAK  
RICHARD V. ZANZALARI\*  
MICHELLE L. BLAT  
PAUL LIM  
ELIZABETH PIETROWSKI

NEW YORK  
DAVIDSON, DAVIDSON & KAPPEL, LLC  
485 SEVENTH AVENUE, 14TH FLOOR  
NEW YORK, NY 10018  
T. 212-736-1940  
F. 212-736-2427  
DDK@DDKPATENT.COM

FRANKFURT  
DAVIDSON, DAVIDSON & KAPPEL EUROPE, LLC  
ARNDTSTRASSE 11  
60325 FRANKFURT AM MAIN, GERMANY  
T. +49 (69) 788 088-0  
F. +49 (69) 788 088-29  
FRANKFURT@DDKPATENT.COM

\*ADMITTED IN NEW JERSEY ONLY  
\*\*DDK EUROPE

September 22, 2004

**VIA PRIORITY MAIL**

Alan Koller, Ph.D., Esq.  
Sr. Assistant General Counsel  
Purdue Pharma LP  
One Stamford Forum  
Stamford, CT 06901

Re: U.S. Patent Application No. 10/056,475  
Entitled: **ANALGESIC COMBINATION OF OXYCODONE  
AND 6-METHOXY-2-NAPHTHYLACETIC ACID**  
Euro-Celtique, S.A.  
Purdue Ref. No.: PT0176  
Our Ref. No.: 200.1079CON3


Dear Alan:

We have now received an Office Action for the above-referenced patent application, a copy of which along with related papers are enclosed for your review.

A response to the Office Action is due **December 7, 2004**, although extensions of time are obtainable if necessary.

Absent your instructions to the contrary, we shall prepare a draft response for your review and consideration prior to the due date. On the other hand, if you have any comments or suggestions concerning this Office Action, we look forward to receiving the same.

Very truly yours,

  
Clifford M. Davidson

CMD:ie  
Enclosure

cc: Robert J. Paradiso, Esq.

**BEST AVAILABLE COPY**



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office  
Address: COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450  
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/056,475	01/25/2002	Ronald M. Burch	200.1079CON3	5119

7590 09/07/2004  
Davidson, Davidson & Kappel, LLC  
14th Floor  
485 Seventh Avenue  
New York, NY 10018

**RECEIVED**  
SEP 09 2004

EXAMINER	
CELSA, BENNETT M	
ART UNIT	PAPER NUMBER
1639	

DATE MAILED: 09/07/2004

DAVIDSON, DAVIDSON & KAPPEL

Please find below and/or attached an Office communication concerning this application or proceeding.

Excel 9-10-04  
IPM 9-10-04

3-7-05 OFFICE ACTION RESPONSE  
Due (Deadline Date)  
12-7-04 OFFICE ACTION RESPONSE  
Due (3 month date)  
11-7-04 Reminder  
10-7-04 send reporting letter

CMD/RSP/RUZ

**Office Action Summary**

SEP 28 2004

Application No.

10/056,475

Applicant(s)

BURCH ET AL.

Examiner

Bennett Celsa

Art Unit

1639

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

**Period for Reply**

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If the period for reply specified above is less than thirty (30) days, a reply within the statutory minimum of thirty (30) days will be considered timely.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133). Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

**Status**

- 1) ☒ Responsive to communication(s) filed on 14 June 2004.
- 2a) ☐ This action is FINAL. 2b) ☒ This action is non-final.
- 3) ☐ Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

**Disposition of Claims**

- 4) ☒ Claim(s) 37-43 and 45 is/are pending in the application.
- 4a) Of the above claim(s) \_\_\_\_\_ is/are withdrawn from consideration.
- 5) ☐ Claim(s) \_\_\_\_\_ is/are allowed.
- 6) ☒ Claim(s) 37-43 and 45 is/are rejected.
- 7) ☐ Claim(s) \_\_\_\_\_ is/are objected to.
- 8) ☐ Claim(s) \_\_\_\_\_ are subject to restriction and/or election requirement.

**Application Papers**

- 9) ☐ The specification is objected to by the Examiner.
- 10) ☐ The drawing(s) filed on \_\_\_\_\_ is/are: a) ☐ accepted or b) ☐ objected to by the Examiner.  
Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).  
Replacement drawing sheet(s) including the correction is required if the drawing(s) is objected to. See 37 CFR 1.121(d).
- 11) ☐ The oath or declaration is objected to by the Examiner. Note the attached Office Action or form PTO-152.

**Priority under 35 U.S.C. § 119**

- 12) ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
- a) ☐ All b) ☐ Some \* c) ☐ None of:
1. ☐ Certified copies of the priority documents have been received.
  2. ☐ Certified copies of the priority documents have been received in Application No. \_\_\_\_\_.
  3. ☐ Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

\* See the attached detailed Office action for a list of the certified copies not received.

**Attachment(s)**

- |                                                                                                    |                                                                             |
|----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| 1) <input checked="" type="checkbox"/> Notice of References Cited (PTO-892)                        | 4) <input type="checkbox"/> Interview Summary (PTO-413)                     |
| 2) <input type="checkbox"/> Notice of Draftsperson's Patent Drawing Review (PTO-948)               | Paper No(s)/Mail Date. _____                                                |
| 3) <input checked="" type="checkbox"/> Information Disclosure Statement(s) (PTO-1449 or PTO/SB/08) | 5) <input type="checkbox"/> Notice of Informal Patent Application (PTO-152) |
| Paper No(s)/Mail Date <u>12/2/02; 1/25/02</u>                                                      | 6) <input type="checkbox"/> Other: _____                                    |

## **DETAILED ACTION**

### ***Status of the Claims***

Claims 37-43 and 45 are currently pending and under consideration..

### ***Election/Restriction***

1. Applicant's election without traverse of Group II (claims 37-43 and 45 methods of treating pain using oxycodone and 6-methoxy-2-naphthylacetic acid or 6-MNA in the correspondence dated 6/14/04 is acknowledged.

### ***Priority***

Applicant should update the cross-reference to parent application which has subsequently issued as a patent.

### ***Claim Rejections - 35 USC § 103***

2. The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

This application currently names joint inventors. In considering patentability of the claims under 35 U.S.C. 103(a), the examiner presumes that the subject matter of the various claims was commonly owned at the time any inventions covered therein were made absent any evidence to the contrary. Applicant is advised of the obligation under 37 CFR 1.56 to point out the inventor and invention dates of each claim that was not commonly owned at the time a later invention was made in order for the examiner to

Art Unit: 1639

consider the applicability of 35 U.S.C. 103<sup>o</sup> and potential 35 U.S.C. 102(e), (f) or (g) prior art under 35 U.S.C. 103(a).

3. Claims 37-40, 42-43 and 45 are rejected under 35 U.S.C. 103(a) as being unpatentable over Baker et al. US Pat. No. 4,569,937 (2/86) and Richardson et al. Drug Safety (Oct. 1996) Vol. 15(4) pages 249-260.

Baker et al. teach pharmaceutical compositions for relieving pain in humans or mammals (e.g. mice, rats etc.) comprising a combination of :

a. a narcotic analgesic (preferably oxycodone: see formulations col. 4-8; mice data in col. 8-10; patent claims), or a pharmaceutically acceptable salt thereof; and  
b. a non-steroidal anti-inflammatory drug or NSAID (preferably ibuprofen: see col. 1-2), or a pharmaceutically acceptable suitable salt thereof,  
in a weight ratio of about 1:800 (e.g. .001:1) to 1:1 (compare to present claim 47: See col. 2)

with oxycodone amounts of about 5 mgs-600mgs (compare to present claim 46).

The Baker reference teaches oral administration (e.g. see present claim 39), which can be coadministered in a "single dosage form" (e.g. see col. 3-8: and present claim 40) or sequentially administered (e.g. as in present claim 42; see i.e. col. 8-9 ; "... mice are dosed sequentially..."). The Baker et al. reference teach that dose ratios can be adjusted and that the analgesic activity of the combined oxycodone and ibuprofen activity is "unexpectedly enhanced" or synergistic "i.e. the resulting activity is greater than the activity expected from the sum of the activities of the individual components", thereby permitting "reduced dosages of narcotic analgesics" (e.g. oxycodone) AND

Art Unit: 1639

which diminishes adverse side effects (e.g. addiction) and toxicity which would result from the otherwise required amounts of the individual drug components" resulting from high dosages of oxycodone or NSAID's such as ibuprofen. See e.g. col. 1-2; col. 3, lines 19-32 (e.g. compare to present 43 and 44 "reduced" active ingredients).

Accordingly, Baker would teach the use of therapeutic and subtherapeutic amounts of oxycodone and/or ibuprofen in view of the synergistic nature of the combinations and the desire to reduce the toxicity and/or side-effects of both agents; and as required by the doctor for his/her particular patient., including dosage optimization e.g. dosage overlapping of active ingredients. See e.g. col. 3 where dosage is modified to suit the particular patient.

The Baker analgesic composition differs from that presently claimed in that it fails to teach the substitution of nimesulide for ibuprofen, or alternatively, the further incorporation of (e.g. encompassed by "consisting essentially of") Nimesulide into the Baker compositions.

Richardson et al.. teach that 6-methoxy-2-naphthylacetic acid (or 6-MNA), the active metabolite of nabumetone, and like nabumetone and Ibuprofen, is more COX-2 than COX-1 selective and as such this non-steroidal anti-inflammatory drug (e.g. NSAID) is associated with a favourable adverse effect profile with low incidence of GI bleeding and low toxicity. E.g. see abstract, page 253 (table 1) and pages 255-256.

Accordingly, one of ordinary skill in the art would have been motivated to substitute 6-MNA (a NSAID) for ibuprofen (a different NSAID) in the Baker reference compositions in light of the Richardson et al. reference teaching that 6-MNA is as

Art Unit: 1639

efficacious, and safe, with few side effects (e.g. as compared to other less selective COX-2 inhibitor NSAID's including ibuprofen).

Alternatively, one of ordinary skill in the art would have been motivated to incorporate 6-MNA, with its potent analgesia and reduced side-effect, into the Baker ibuprofen/oxycodone compositions in order to reduce the amounts (e.g. therapeutic/subtherapeutic) of ibuprofen/oxycodone in order to further avoid the side effects (e.g. addiction) or toxicity resulting from ibuprofen/oxycodone.

Additionally, it is noted that the instant situation is amenable to the type of analysis set forth in *In re Kerkhoven*, 205 USPQ 1069 (CCPA 1980) wherein the court held that it is *prima facie* obvious to combine two (or more) compositions each of which is taught by the prior art to be useful for the same purpose

Thus, it would have been *prima facie* obvious to one of ordinary skill in the art at the time of applicant's invention to modify the Baker reference analgesic composition by substituting the NSAID Nimesulide (for the NSAID ibuprofen) or supplement Baker's composition with 6-MNA in light of the benefits of 6-MNA increased safety/decreased side effect as compared to other NSAID's as taught by the Richardson et al. reference.

4. Claims 37-43 and 45 are rejected under 35 U.S.C. 103(a) as being unpatentable over the Baker et al. '937 and Richardson et al. applied to claims 37-40, 42-43 and 45 above, and further in view of Mayer et al. US Pat. No. 5834,479 (11/98).

The teaching of the Baker and Richardson et al. references recited above is hereby incorporated by reference in its entirety.

To the extent that the Baker and Richardson et al. references fail to teach the administration of the analgesia active agent (e.g. 6-MNA) "before, ... with, or after" administration of the oxycodone" (particularly before/after ) (e.g. see present claim 41) the Mayer et al. reference is cited.

The Mayer et al. reference teaches that analgesia effectiveness of an analgesia active agent (e.g. a NSAID, such as ibuprofen see i.e. table in col. 7) can be "significantly enhanced" by administering (e.g. oral administration) the active agent "prior to, with or following the administration of an analgesia enhancer" (e.g. a nontoxic NMDA receptor blocker and/or a nontoxic substance that blocks at least one major intracellular consequence of NMDA receptor activation) such as "dextromethorphan", which is the D-isomer of codeine . See e.g. col. 1; patent claims.

Accordingly, the Mayer et al. reference provides motivation to one of ordinary skill in the art to not only co- administer different analgesic agents to achieve enhanced analgesia, but to also administer the NSAID prior or subsequent to the second analgesic agent i.e. an analgesia enhancer, which includes codeine or its derivatives (e.g. dextromethorphan, dextrorphan, oxycodone etc.) .

Thus, it would have been prima facie obvious to one of ordinary skill in the art at the time of applicant's invention to modify the combined Baker and Richardson et al. reference teachings by administering one of the analgesic active agents (e.g. 6-MNA) "before, ... with, or after" administration of the second analgesic agent (e.g. oxycodone) in order to obtain significantly enhanced analgesia.



Art Unit: 1639


**Conclusion**

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Bennett Celsa whose telephone number is 571-272-0807. The examiner can normally be reached on 8-5.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Andrew Wang can be reached on 571-273-0811. The fax phone number for the organization where this application or proceeding is assigned is 703-872-9306.

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see <http://pair-direct.uspto.gov>. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free).

Bennett Celsa  
Primary Examiner  
Art Unit 1639



BC  
August 26, 2004

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1078CON3

SERIAL NO.: Not yet known

10/056,475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Herewith

GROUP: Not Yet Known

1639

U.S. PAT. & TM. OFF.  
10/056475  
01/25/02

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER							DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
<i>[Signature]</i>	AA	5	8	5	8	2	5	7	01/12/99	Talley	548	247	08/14/88
	AB	5	8	8	3	8	2	2	01/26/99	Mayer, et al.	514	270	07/02/99
	AC	5	8	4	0	7	3	1	11/24/98	Mayer, et al.	514	289	08/02/95
	AD	5	8	6	8	4	8	8	02/09/99	Mayer, et al.	514	282	07/07/97
	AE	5	8	6	1	4	1	8	01/19/99	Dube, et al.	514	334	07/11/97
	AF	5	7	8	9	4	1	3	08/04/98	Black, et al.	514	255	01/21/97
	AG	5	6	0	4	2	6	0	02/18/97	Guay, et al.	514	605	11/04/93
	AH	5	4	5	8	8	7	8	10/17/95	Singh, et al.	424	400	09/30/94
	AI	5	5	1	6	8	0	3	05/14/96	Raffa	514	570	03/01/85
<i>[Signature]</i>	AJ	5	8	4	3	4	6	8	12/01/98	Burkoth, et al.	424	448	05/13/96

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

<i>[Signature]</i>	AK	Differential Inhibition of Cyclooxygenase-1 (COX-1) and -2 (COX-2) By NSAIDs: Consequences on Anti-Inflammatory Activity Versus Gastric and Renal Safety, M. Pairet, et al., Inflammopharmacology 4: 61-70, (1996).
	AL	Differential effects of inhibitors of cyclooxygenase (cyclooxygenase 1 and cyclooxygenase 2) in acute inflammation, Derek W. Gilroy, et al. European J. Pharm 355 pp 211-217, (1998).
	AM	Cyclooxygenases 1 and 2, J.R. Vane, et al. Annu. Rev. Pharmacol. Toxicol. 38: 97-121, (1998).
	AN	Analysis of the effects of cyclooxygenase (COX)-1 and COX-2 in spinal nociceptive transmission using indomethacin, a non-selective COX inhibitor, and NS-398, a COX-2 selective inhibitor, Tatsuo Yamamoto, et al. Brain Research 739:104-110, (1996).
	AO	Comparative Analgesic Efficacy of Nimesulide and Diclofenac Gels after Topical Application on the Skin, S. Sengupta, et al., Skin Pharmacol. And Applied Skin Phys. 11:273-278, (1998).
	AP	Carrageenan-induced hyperalgesia is associated with increased cyclo-oxygenase-2 expression in spinal cord, Cartiona Hay and Jacqueline de Bellerche, Neuro Report 8, 1249-1251, (1997).
	AQ	The Mechanisms of Action of NSAIDs in Analgesia, Jeremy N. Cashman, Drugs 52 Supp. 5:13-23, (1996).
	AR	Differential effects of inhibition of isoforms of cyclooxygenase (COX-1, COX-2) in chronic inflammation, D.W. Gilroy, et al. Inflamm. Res. 47:79-85, (1998).
<i>[Signature]</i>	AS	Constitutive Cyclooxygenase (COX-1) and Inducible Cyclooxygenase (COX-2): Rationale for Selective Inhibition and Progress to Date, Don E. Griswold and Jerry L. Adams, Medicinal Research Reviews, Vol. 16, No. 2, pp. 181-206, (1996).

EXAMINER

DATE CONSIDERED

8/26/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1079CON3

SERIAL NO.: Not Yet Known

10/056, 275

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known

GROUP: Not Yet Known

1639

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	BA						
	BB						
	BC						
	BD						
	BE						
	BF						
	BG						
	BH						
	BI						
	BJ						
	BK						

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
							YES	NO
	BL							

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	BM	Cyclooxygenase in biology and disease, Raymond N. Dubois, et al., FASEB J. Vol. 12 pp. 1063-1073 (1998).
	BN	Pharmacology of Meloxicam, A new Non-Steroidal Anti-Inflammatory Drug With An Improved Safety Profile Through Preferential Inhibition of COX-2, G. Engelhardt, British J. Rheumatology, 35 (supp 1):4-12, (1996)
	BO	Cyclooxygenase 1 Contributes to Inflammatory Responses in Rats and Mice; Implications for Gastrointestinal Toxicity. John L. Wallace, et al. Gastroenterology, 115:101-109 (1998).
	BP	Distinct isoforms (COX-1 and COX-2) of cyclooxygenase: possible physiological and therapeutic implications. M. Pairet and G. Engelhardt, Fundam. Clin. Pharmacol. 10:1-15, (1996).
	BQ	Involvement of Prostaglandins Produced by Cyclooxygenase-1 in Murine Visceronoception Induced by Phenylquinone, Hidenobu Kusuha, et al. Prostaglandins 55: 43-49, (1998).
	BR	Effect of COX-1 and COX-2 Inhibition on Induction and Maintenance of Carrageenan-Evoked Thermal Hyperalgesia in Rats. D. Dirig, et al. J. Pharmacol. And Experimental Therapeutics Vol. 285, No. 3. pp 1031-1038.

EXAMINER

DATE CONSIDERED

10/5/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICEATTY. DOCKET NO.:  
200.1079CON3

SERIAL NO.: Not Yet Known

10/056, 475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known

GROUP: Not Yet Known

1639

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	CA						
	CB						
	CC						
	CD						

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
							YES	NO
	CE	9 9 0 7 4 1 3	2/18/99	WO				
	CF	9 9 3 2 1 1 9	7/1/99	WO				
	CG							
	CH							
	CI							
	CJ							

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	CK	Effect of meloxicam on postoperative pain after abdominal hysterectomy, J.P. Thompson et al. British Journal of Anaesthesia 84 (2) 151-4 (2000).
	CL	Intrathecal cyclooxygenase inhibitor administration attenuates morphine antinociceptive tolerance in rats. C.S. Wong et al., British Journal of Anaesthesia 85 (5) 747-51 (2000).
	CM	Cyclooxygenase inhibitors increase morphine effects on mesolimbic dopamine neurons, M. Melis, et al. Eur. J. Pharmacology 387 (1) R1-R3 (2000).
	CN	Synergistic antiallodynic effects of spinal morphine with ketorolac and selective COX-1 and COX-2 inhibitors in nerve-injured rats, J.M. Lashbrook, et al. Pain 82 (1) 65-72 (1999).
	CO	Enhancement of opioid inhibition of gaba-ergic synaptic transmission by cyclo-oxygenase inhibitors in rat periaqueductal grey neurones, Vaughn et al. British Journal of Pharmacology 123 (8) 1479-81 (1998).

EXAMINER

DATE CONSIDERED

8/24/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1079CON3

SERIAL NO.: Not Yet Known  
10/056,475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known

GROUP: Not Yet Known  
1639

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER							DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
<i>[Signature]</i>	DA	5	4	7	4	9	9	5	12/12/95	Ducharme et al.	514	241	01/10/94
	DB	5	6	9	1	3	7	4	11/25/97	Black et al.	514	473	05/18/95
	DC												
	DD												
	DE												
	DF												
	DG												
	DH												
	DI												
	DJ												
	DK												

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER							DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
													YES	NO
	DL													
	DM													
	DN													
	DO													
	DP													

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	DO	
	DR	
	DS	

EXAMINER

*[Signature]*

DATE CONSIDERED

8/20/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 608; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1078CON3

SERIAL NO.: Not Yet Known

10/054, 475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)



APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known




GROUP: Not Yet Known

1639



## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER							DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	EA	3	8	0	0	0	4	1	03/26/74	Müller, et al.	424	273	03/02/71
	EB	4	3	2	2	4	2	7	03/30/82	Buyinski, et al.	424	260	04/16/81
	EC	4	3	3	8	3	2	4	06/06/82	Gardocki	424	266	03/17/81
	ED	4	4	0	4	2	1	0	09/13/83	Schmidt	424	260	06/30/82
	EE	4	4	0	7	8	0	4	10/04/83	Schmidt	424	260	06/30/82
	EF	4	4	0	7	8	0	5	10/04/83	Schmidt	424	260	08/30/82
	EG	4	4	6	4	3	7	6	08/07/84	Sunshine, et al.	424	253	10/11/83
	EH	4	4	8	6	4	3	6	12/04/84	Sunshine, et al.	424	253	03/11/83
	EI	4	4	8	9	0	8	0	12/18/84	Lomen	424	260	12/06/82

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER							DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
													YES	NO
	EJ	2	0	2	3	3	4	9	02/18/91	Canada	A61K	31/44		
	EK	0	2	7	4	8	4	5	07/20/88	EP (A1)	A61K	31/19		
	EL	0	3	8	8	1	2	5	09/19/90	EP (A1)	A61K	31/485		

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	EM	Pharmacokinetics and Drug Input Characteristics for a Diclofenac-Codeine Phosphate Combination Following Oral and Rectal Administration A. Hansen, et al. <i>Arzneim-Forsch./Drug Res.</i> 46 (II) 57-63 (1996).
	EN	Comparison of a Standard Ibuprofen Treatment Regimen with a New Ibuprofen/Paracetamol/Codeine Combination in Chronic Osteo-arthritis G.K. Vlok, et al. Univ. Stellenbosch and Tygerberg Hospital, Dept. Orthopaedic Surg. pp 3-6, (1987).
	EO	
	EP	
	EQ	
	ER	

EXAMINER

DATE CONSIDERED

8/26/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1078CON3

SERIAL NO.: Not Yet Known

10/054,475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known

GROUP: Not Yet Known

1439

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER							DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
<i>MM</i>	FA	4	5	6	7	1	8	3	01/28/86	Sunshine, et al.	514	264	12/12/83
	FB	4	5	6	9	9	3	7	02/11/86	Baker, et al.	514	282	02/11/83
	FC	4	5	7	1	4	0	0	02/18/86	Arnold	514	282	12/18/84
	FD	4	5	8	7	2	5	2	05/06/86	Arnold	514	282	12/18/84
	FE	4	6	1	9	9	3	4	12/28/86	Sunshine, et al.	514	277	07/08/85
	FF	4	6	9	0	9	2	7	09/01/87	Voss, et al.	514	282	02/03/86
	FG	4	8	3	9	1	7	6	06/13/89	Pakhania, et al.	424	465	12/08/87
	FH	4	8	4	4	9	0	7	07/04/89	Elger, et al.	424	465	08/14/86
	FI	4	9	2	7	8	5	4	05/22/90	Sunshine, et al.	514	570	08/21/89
	FJ	5	1	6	4	3	9	8	12/17/92	Sims, et al.	514	282	04/01/91
	FK	5	1	9	0	9	4	7	03/02/93	Riess, et al.	514	282	08/16/91
<i>MM</i>	FL	5	2	4	0	6	9	4	08/31/93	Gwaltney, Jr.	424	45	12/19/91

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER							DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
													YES	NO
<i>MM</i>	FM	0	0	6	8	8	3	8	06/25/82	EP (A1)	A61K	31/485		
	FN	0	0	6	8	8	3	8	06/25/82	EP (B1)	A61K	31/485		
<i>MM</i>	FO	2	0	8	1	6	0	4	05/01/95	Canada	A61K	031/135		
	FP													
	FQ													

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	FR	
	FS	
	FT	

EXAMINER

*MM*

DATE CONSIDERED

8/26/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.: 200.1079CON3

SERIAL NO.: Not Yet Known  
101056, 475

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: Not Yet Known

GROUP: Not Yet Known  
1639

## U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
MA	GA	5 4 0 9 8 4 4	04/25/95	Black, et al.	514	359	03/12/93
	GB	5 4 3 6 2 6 5	06/25/95	Black, et al.	514	420	11/12/93
	GC	5 5 1 0 3 6 8	04/23/96	Lau, et al.	514	419	05/22/95
	GD	5 5 2 1 2 1 3	05/28/96	Prasit, et al.	514	443	08/28/94
	GE	5 5 5 0 1 4 2	08/27/96	Ducharme et al.	514	365	05/08/95
	GF	5 5 5 2 4 2 2	09/03/96	Gauthier, et al.	514	368	01/11/95
	GG	5 6 0 4 2 5 3	02/18/97	Lau, et al.	514	415	05/22/95
	GH	5 6 3 9 7 8 0	06/17/97	Lau, et al.	514	419	05/22/95

## FOREIGN PATENT DOCUMENTS

	DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
						YES	NO
GI							
GJ							
GK							
GL							
GM							

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

GN	Anti-inflammatory Drugs and Their Mechanism of Action J.R. Vane, et al. Inflamm. Res. 47, Supplement 2 (1998).
GO	
GP	
GQ	
GR	

EXAMINER

DATE CONSIDERED

8/26/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.



[illegible]

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE:  
September 17, 1998GROUP: 1627  
639

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER							DATE	COUNTRY	CLASS	SUBCLAS S	TRANSLATION	
													YES	NO
	AA	0	6	5	4	2	6	3	05/24/95	EP (A1)	A61K	31/135		
	AB	0	6	5	4	2	6	3	05/24/95	EP (B1)	A61K	31/135		
	AC	9	7	1	7	9	7	8	05/22/97	WO	A61K	33/00		
	AD	9	7	2	5	9	8	8	07/24/97	WO	A61K	31/495		
	AE	9	7	3	2	8	5	7	09/12/97	WO	C07D	241/104		

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

AF	Dray et al. New Pharmacological Strategies for Pain Relief. <u>Annual Review of Pharmacology &amp; Toxicology</u> , 36, pp. 253-280. (1996).
AG	Brasseur, L. Revue des therapeutiques pharmacologiques actuelles de la douleur. <u>Drugs</u> , 53 Suppl 2, pp. 10-17. (1997)
AH	Rang et al. New molecules in analgesia. <u>British Journal of Anesthesia</u> , 75, pp. 145-156 (1995)
AI	Beaver, WT. Combination Analgesics. <u>American Journal of Medicine</u> , 77 (Suppl 3A), pp. 38-53. (1984).
AJ	Beaver, WT. Chapter 29: Nonsteroidal Antiinflammatory Analgesics and Their Combinations with Opioids. In <u>Evaluation and Treatment of Chronic Pain</u> , 2 <sup>nd</sup> ed., William & Wilkins pp. 363-383. (1992).
AK	Goodman & Gilman's. <u>The Pharmacological Basis of Therapeutics</u> , 9 <sup>th</sup> Edition. McGraw-Hill, New York, pp 535 and 551-552.
AL	Picard et al. Ketorolac potentiates morphine in postoperative patient-controlled analgesia. <u>Pain</u> , 73, 3 pp. 401-406. (1997).
AM	Etches et al. Continuous Intravenous Administration of Ketorolac Reduces Pain and Morphine Consumption After Total Hip or Knee Arthroplasty. <u>Anesthesia &amp; Analgesia</u> , 81 (6), pp. 1175-1180. (1995).
AN	Hodsman et al. The morphine sparing effects of diclofenac sodium following abdominal surgery. <u>Anaesthesia</u> , 42(9), pp. 1005-1008. (1987).
AO	Kaasalainen et al. Developments in the treatment of cancer pain in Finland: The third nation-wide survey. <u>Pain</u> , 70, 2-3, pp. 175-183. (1997).
AP	Sunshine et al. Analgesic Efficacy of a Hydrocodone with Ibuprofen Combination Compared with Ibuprofen Alone for the Treatment of Acute Postoperative Pain. <u>Journal of Clinical Pharmacology</u> , 37 (10), pp. 908-915. (1997).
AQ	Insel. Chapter 27: Analgesic-Antipyretic and Anti-Inflammatory Agents. In Hardman, ed., <u>Goodman &amp; Gilman's The Pharmacological Basis of Therapeutics</u> , 9 <sup>th</sup> Edition. McGraw-Hill, New York, pp. 654-655. (1996).
AR	Pollsson. Non-steroidal Anti-inflammatory Drugs: Practical and Theoretical Consideration in Their Selection. <u>The American Journal of Medicine</u> , 100 (Suppl 2A), pp. 2A-31S - 2A-36S. (1996).
AS	Vane, J. Towards a better aspirin. <u>Nature</u> , 367, pp. 215-216. (1994).
AT	Simon, L.S. Nonsteroidal Antiinflammatory Drugs and Their Effects: The Importance of COX 'Selectivity'. <u>Journal of Clinical Rheumatology</u> , 2 (3), pp. 135-140. (1996).

EXAMINER

DATE CONSIDERED

8/26/98

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

10/054,277

## LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE: September  
17, 1998

GROUP: 1627

1439

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

BA	Van Ryn et al. Selective cyclooxygenase-2 inhibitors: pharmacology, clinical effects, and therapeutic potential. <u>Expert Opinion On Investigational Drugs</u> , pp. 609-614. (1997).
BB	Vane et al. New insights into the mode of action of anti-inflammatory drugs. <u>Inflammation Research</u> , 44, (No.1), pp 1-10 (1995).
BC	Engelhardt. Meloxicam: A Preferential Inhibitor of COX-2. <u>British Journal of Rheumatology</u> , 34, Abstract Suppl. 1, p. 48. (1995). Abstract.
BD	Lane, N.E. Pain Management in Osteoarthritis: The Role of COX-2 Inhibitors. <u>Journal of Rheumatology</u> , Vol. 24, Suppl 49, pp. 20-24. (1997).
BE	Boyce et al. L-745,337: A Selective Inhibitor of Cyclooxygenase-2 Elicits Antinociception But Not Gastric Ulceration in Rats. <u>Neuropharmacology</u> Vol. 33, pp. 1609-1611. (1994).
BF	Donnelly et al. COX-II Inhibitors - a new generation of safer NSAIDS? <u>Alimentary Pharmacology and Therapeutics</u> , 11, 2, pp. 227-236. (1997).
BG	Wallace, J.L. Nonsteroidal Anti-inflammatory Drugs and Gastroenteropathy: The Second Hundred Years. <u>Gastroenterology</u> , 112, 3, pp. 1000-1016. (1997).
BH	Robinson, D.R. Regulation of Prostaglandin Synthesis by Antiinflammatory Drugs. <u>J Rheumatology</u> , 24, Suppl. 47, pp. 32-39. (1997).
BI	Tannenbaum et al. An Evidence-Based Approach to Prescribing NSAIDS in Musculoskeletal Disease: A Canadian Consensus. <u>Canadian Medical Association Journal</u> , 155, 1, pp. 77-88. (1996).
BJ	Mehlisch et al. Analgesic Efficacy and Plasma Levels of a Highly Selective Inhibitor of COX-2 (SC-58635, SC) in Patients with Postsurgical Dental Pain. <u>Journal of Clinical Pharmacology</u> , 37, 9, 863. (1997). Abstract.
BK	Dammann. Selective COX-2 Inhibition: Its Relevance for NSAID-Gastrointestinal Toxicity. <u>Gut</u> , 39, Suppl. 3, A151. (1996). Abstract.
BL	Penning et al. Synthesis and Biological Evaluation of the 1, 5 -diarylpyrazole class of cyclooxygenase-2 inhibitors: Identification of 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide (SC-58635, Celecoxib). <u>Journal Of Medicinal Chemistry</u> , 40(9), 1347-65. (1997).
BM	Lipsky et al. Outcome of Specific COX-2 Inhibition in Rheumatoid Arthritis. <u>Journal Of Rheumatology</u> , 24 Suppl 49, pp. 9-14. (1997).
BN	
BO	
BP	
BQ	
BR	
BS	
BT	

EXAMINER

DATE CONSIDERED

8/24/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449  
(REV. 7-80)

U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

ATTY. DOCKET NO.:  
200.1079US

SERIAL NO.:  
09/154,354 10/054,475

LIST OF PRIOR ART CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT(S): Ronald M. BURCH, et al.

FILING DATE:  
September 17, 1998

GROUP:  
1627 1439

U.S. PATENT DOCUMENTS

*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	AA						
	AB						
	AC						
	AD						
	AE						
	AF						
	AG						

FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
							YES	NO
	AH	0 7 3 4 2 7 5	11/04/1997	Australia	C07D	261/08		
	AI							
	AJ							
	AK							
	AL							

OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	AM	
	AN	
	AO	
	AP	
	AQ	
	AR	
	AS	

EXAMINER

DATE CONSIDERED

8/26/04

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

<b>Notice of References Cited</b>	Application/Control No. 10/056,475	Applicant(s)/Patent Under Reexamination BURCH ET AL.	
	Examiner Bennett Celsa	Art Unit 1639	Page 1 of 1

**U.S. PATENT DOCUMENTS**

*		Document Number Country Code-Number-Kind Code	Date MM-YYYY	Name	Classification
	A	US-5,834,479	11-1998	Mayer et al.	514/289
	B	US-			
	C	US-			
	D	US-			
	E	US-			
	F	US-			
	G	US-			
	H	US-			
	I	US-			
	J	US-			
	K	US-			
	L	US-			
	M	US-			

**FOREIGN PATENT DOCUMENTS**

*		Document Number Country Code-Number-Kind Code	Date MM-YYYY	Country	Name	Classification
	N					
	O					
	P					
	Q					
	R					
	S					
	T					

**NON-PATENT DOCUMENTS**

*		Include as applicable: Author, Title Date, Publisher, Edition or Volume, Pertinent Pages)
	U	Richardson et al. Drug Safety (Oct. 1996) Vol. 15(4) pages 249-260.
	V	
	W	
	X	

\*A copy of this reference is not being furnished with this Office action. (See MPEP § 707.05(a).)  
Dates in MM-YYYY format are publication dates. Classifications may be US or foreign.

# The Clinical Implications of Inhibition of the Inducible Form of Cyclo-Oxygenase

Charles Richardson and Paul Emery

Department of Rheumatology and Rehabilitation, University of Leeds, Leeds, England

## Summary

There are 2 isoenzymes of cyclo-oxygenase (COX). There is a constitutive enzyme COX-1 which has a wide tissue distribution. In addition there is an inducible enzyme COX-2 which has a restricted tissue distribution. The inducible enzyme COX-2 is responsible for the generation of prostaglandins at sites of tissue inflammation and its inhibition is associated with an anti-inflammatory action. The constitutive enzyme, COX-1 is responsible for the production of prostaglandins with multiple functions. One well known and clinically important function is the gastroprotective effect of the prostaglandins produced in the gastric mucosa. While selective COX-2 inhibition may be associated with a reduced incidence of gastric adverse effects, a concern remains that its inhibition at other locations may be associated with other adverse effects.

Inflammation is the body's response to tissue damage. The major purpose of this immensely complex response is to permit the entry of fluid, proteins and cells from the bloodstream into the damaged tissues to produce tissue healing. The major features of inflammation are: vasodilatation, increased vascular permeability and cellular infiltration. The net effect is an increased blood supply and entry of proteins and cells into the affected site. In clinical practice this produces pain, swelling and erythema. One of the major groups of chemical mediators involved in the inflammatory response is the prostaglandins, which are synthesised by the enzyme cyclo-oxygenase (COX) from arachidonic acid.<sup>[1]</sup> It is now apparent that there are 2 isoenzymes of COX, a constitutive enzyme COX-1 responsible for the production of prostaglandins with 'general housekeeping' functions in the body, such as maintenance of renal perfusion and a protective effect in the gastric mucosa against ulcer-

ation, and an inducible enzyme COX-2 responsible for the production of the pro-inflammatory prostaglandins. The nonsteroidal anti-inflammatory drugs (NSAIDs), by inhibiting the COX enzyme, are able to reduce the pain and swelling of inflammation. In theory, if new molecules that preferentially inhibit the inducible enzyme alone could be developed then these would have an anti-inflammatory effect and would not affect the production of prostaglandins for general 'housekeeping functions'. In this article we will review the clinical experience of COX selectivity as a determinant of NSAID toxicity.

## 1. Prostaglandin Metabolism

The prostaglandins are a family of lipid soluble molecules synthesised by the COX enzyme from arachidonic acid, a 20 carbon unsaturated fatty acid.<sup>[2]</sup> There are at least 16 different prostaglandins and these can be divided into 9 chemical




Fig. 1. The pathways of prostaglandin biosynthesis. Abbreviation: PG = prostaglandin.

classes. The prostaglandins do not exist free in the tissues but are synthesised and released in response to the appropriate stimulus. Their biosynthetic pathways are illustrated in figure 1. Prostaglandins are found in many human tissues and have a diverse range of effects.

For immunoregulation prostaglandin (PG) E<sub>2</sub> has been most extensively studied and has both pro-inflammatory and anti-inflammatory roles. It is found in high levels in acutely inflamed tissue where it is responsible for vasodilatation<sup>[3]</sup> and sensitises nerve endings to bradykinin and histamine, resulting in the pain characteristic of inflammation (i.e. pro-inflammatory action).<sup>[4]</sup> In chronic inflammation PGE<sub>2</sub>, again produced by COX-2,<sup>[5]</sup> has an anti-inflammatory action and suppresses a wide variety of cellular functions including lymphocyte proliferation, T cell cytotoxicity and leucocyte motility. A further important role of PGE<sub>2</sub> [and prostacyclin (PGI<sub>2</sub>)] is as a vasodilator to maintain renal perfusion; this is of particular importance at times of circulatory collapse.<sup>[6]</sup> PGI<sub>2</sub> has a lesser role in inflammation but a major one as a gastroprotective agent.<sup>[7]</sup>

In addition, prostaglandins have an important role in haemostasis and inhibition of their production results in a bleeding tendency.<sup>[8]</sup> Therefore the blanket inhibition of COX activity by NSAIDs results in an anti-inflammatory action, a tendency to peptic ulceration and renal toxicity.

It is now clear that there are 2 isoenzymes of COX: there is a constitutive enzyme (COX-1) with a wide tissue distribution and an inducible enzyme (COX-2).<sup>[9-11]</sup> The constitutive enzyme (COX-1) is responsible for the production of prostaglandins with general (i.e. continuously required) 'house-keeping' functions.<sup>[9]</sup> By contrast, the inducible enzyme has a limited constitutive expression, but is generated in many tissues as a response to an external stimulus. The range of prostaglandins produced by COX-1 and 2 are similar, their physiological role being dependent on their tissue of origin and the stimulus determining their production. For example PGE<sub>2</sub> is synthesised by:

- COX-1 in the gastric mucosa and has a protective effect against the development of mucosal ulceration;

- COX-1 and COX-2 in the renal mesangium and has a role in renal haemostasis and salt balance;
- COX-2 at sites of both acute and chronic tissue inflammation and contributes to pain and vasodilatation.

The identification of COX-2 has produced interest in the development of molecules which preferentially inhibit this inducible isoenzyme alone. In theory such a molecule would have an (improved) anti-inflammatory effect with an improved safety profile.<sup>[12]</sup> However, as stated above it is now becoming increasingly recognised that prostaglandins produced by COX-2 have many other important physiological roles in addition to their pro-inflammatory effect. For example COX-2 has been demonstrated to be responsible for the production of prostaglandins in the kidney, brain, seminal vesicle and uterus, and clearly inhibition of their synthesis at these sites may not be desirable.

## 2. Isoenzymes of Cyclo-Oxygenase (COX)

The 2 isoenzymes of COX have 60% amino acid homology and both metabolise arachidonic acid to make prostaglandins. They have similar binding sites (and binding constants) for arachidonic acid but there are important differences in structure and function: COX-1 has a greater specificity for C<sub>20:4</sub> fatty acids, whereas COX-2 can use 18 or 20 carbon acids equally well and is therefore able to accept a wider range of substrate.<sup>[13]</sup> The isoenzymes are coded for by different genes. The COX-2 gene is larger and codes for 4kb of mRNA, the smaller COX-1 gene codes for 2.8kb of mRNA.<sup>[9,11,14]</sup> Furthermore, expression of the COX-2 gene (but not COX-1) can be modulated by the binding of glucocorticoids,<sup>[10]</sup> interleukin-1, interleukin-6 and phorbol esters.<sup>[15,16]</sup> Therefore the synthesis of COX-2 gene is responsive to a pro-inflammatory stimulus. The clinical implication of COX-2 inhibition can only be understood with a knowledge of the biological effects of prostaglandins produced by this enzyme.

## 3. Biological Effects of Prostaglandins

### 3.1 Role in the Gastric Mucosa

In the gastric mucosa the following physiological roles have been identified for the prostaglandins generated by COX-1 (see reference 17) for a review of this topic):

- Inhibition of basal cell gastric acid production.
- Stimulation of basal cell bicarbonate secretion.
- Stimulation of basal cell gastric mucus secretion. The increase in mucus thickness increases the pH gradient between the lumen and the epithelial surface.
- Enhanced gastric blood flow during acid production, thus preventing the development of erosions that occur when such vasodilatation does not occur.
- Stimulation of rapid repair of disrupted epithelium. This may involve migration of basal cells toward the lumen to repair mucosal injury and enhanced RNA and protein synthesis.

### 3.2 Role in Haemostasis

The enzyme thromboxane synthetase is responsible for the production of thromboxane within platelets, while COX is responsible for the production of PGI<sub>2</sub> in the vascular endothelium. Thromboxane stimulates platelet aggregation which is inhibited by PGI<sub>2</sub>. NSAIDs inhibit both COX and thromboxane synthetase and decrease the production of PGI<sub>2</sub> and thromboxane respectively. It is the balance of inhibition of these 2 enzymes that determines the overall effect a given NSAID has on haemostasis.<sup>[18]</sup> Since the vascular endothelium is capable of synthesising more COX but platelets are not able to synthesise more thromboxane synthetase, treatment with NSAIDs tends to result in a relative increase in PGI<sub>2</sub> levels and a bleeding tendency.

The human endothelium possesses both the constitutive COX-1 and the inducible isoenzyme COX-2.<sup>[18]</sup> Interleukin-1 and 6 have been demonstrated to regulate the transcription of the COX-2 gene in the endothelium.<sup>[18,19]</sup> In addition, activation



of the gene for COX-2 represents an early response of the vascular smooth muscle to injury. Platelet derived growth factor, epidermal growth factor and thrombin are strong inducers whilst interleukin-1 $\alpha$  and acidic and basic fibroblast growth factors are weak inducers.<sup>[20]</sup>

### 3.3 Role in Tissue Inflammation

Under normal conditions resting neutrophils, macrophages and fibroblasts show little or no COX activity and hence prostaglandin production.<sup>[5]</sup> However in response to an inflammatory stimulus (e.g. pro-inflammatory cytokines, bacterial lipopolysaccharide and growth factors) there is increased COX activity due to new gene expression of COX-2. Furthermore fibroblasts and endothelial cells from human rheumatoid synovial tissue synthesise increased amounts of COX-2 in response to stimulation by interleukin-1 and phorbol esters. Glucocorticoids and possibly some NSAIDs can inhibit these effects on COX-2 production.<sup>[15,16]</sup>

### 3.4 Role in Renal Function

It is widely accepted that renal prostaglandins play an important role in the maintenance of renal haemostasis and tubular function. The effects of prostaglandins on the kidney may be summarised as follows.

- Maintenance of normal renal blood flow. PGE<sub>2</sub>, PGF<sub>2 $\alpha$</sub>  and PGI<sub>2</sub> are all potent vasodilators.
- Decreased renal response to antidiuretic hormone (ADH). Prostaglandins inhibit the effect of ADH on the collecting duct.
- Decreased resorption of sodium and chloride from the ascending limb of the loop of Henle in response to PGE<sub>2</sub>.
- Increased renin release. PGE<sub>2</sub> and PGI<sub>2</sub> stimulate the release of renin from the juxtaglomerular apparatus.

It is becoming increasingly recognised that the maintenance of normal renal haemostasis is dependant on the ability to generate prostaglandins in response to an appropriate stimulus. *In vitro* studies have demonstrated that continuous stretching of rat glomerular mesangial cells results in cell prolifer-

ation, protein synthesis and the generation of prostaglandins by COX-2.<sup>[21]</sup> The ability to respond to mechanical stress by the production of vasoactive prostaglandins is thought to be important in renal homeostasis.

Cyclosporin, an immunosuppressive drug with considerable renal toxicity, inhibits the synthesis of COX-2 in rat renal mesangial cells in response to pro-inflammatory cytokines. This may contribute to its nephrotoxicity.<sup>[22]</sup>

### 3.5 Role in Brain Signalling

Prostaglandins have been identified as important neurotransmitters in the brain. Animal experimentation has shown that the COX-2 isoenzyme is widely distributed throughout the forebrain in discrete populations of neurons and is enriched in the cortex and hippocampus.<sup>[23]</sup> Basal expression of COX-2 appears to be regulated by background synaptic activity and can rapidly and transiently induce increased synaptic activity or seizures.<sup>[23,24]</sup> It is likely that COX-2 induction is an important factor in the regulation of cerebral prostaglandin synaptic transmission.

### 3.6 Role in Skeletal Muscle

It has been reported that differentiated skeletal muscle in tissue culture developed a 3-fold increase in COX-2 activity and anabolic myofibre growth in response to repetitive mechanical stimulation. The inhibition of COX-2 expression prevented the muscle fibre growth. The obvious conclusion made was that muscle cell growth was at least in part dependent on COX-2 induction.<sup>[25]</sup>

### 3.7 Role in the Uterus

Prostaglandins are required for normal uterine function. It has been postulated that the induction of COX-2 by uterine stromal cells may be required to support the prostaglandin production necessary for the early stages of embryo implantation.<sup>[26]</sup> In addition, the induction of COX-2 probably mediates the increase in prostaglandin synthesis occurring

within the amniotic membranes of the placenta that occurs at the onset of labour.<sup>[27]</sup>

### 3.8 Role of COX in the Seminal Vesicle

COX-2 was originally described in the seminal vesicle. It has a role in the production of prostaglandins present in semen.<sup>[28]</sup>

## 4. Clinical Significance of COX-2 Inhibition

In addressing the clinical significance of COX-2 inhibition there are 2 questions to be answered. Is selective inhibition of COX-2 associated with a decreased incidence of adverse effects and does selective COX-2 inhibition create its own adverse effects?

### 4.1 COX Selectivity of the Currently Available NSAIDs

In assessing the clinical implications of COX-2 inhibition it is useful first to consider the NSAIDs currently available on the market and determine to what extent their COX selectivity influences their adverse effect profile. The ability of different NSAIDs to inhibit the 2 isoenzymes of COX has been investigated using a number of biological assays.<sup>[26,29-31]</sup> The ability of a molecule to inhibit

COX is determined in terms of its  $IC_{50}$ , the concentration of the compound required to inhibit 50% of COX activity. To compare the activity of NSAIDs on COX-1 and COX-2 the ratio of the  $IC_{50}$  of COX-2 : COX-1 is employed. The lower the ratio, the greater the specificity for COX-2 i.e. a ratio of >1 indicates that a given compound has more activity on COX-2 than COX-1. It should be noted that the different values for COX ratios given by different investigators are due to the use of different COX preparations and that no preparation has been accepted as a 'gold standard'. The results are summarised in table 1. These biological assays demonstrate clear cut differences in the ability of NSAIDs to inhibit the 2 isoenzymes. In the following section the COX selectivity of NSAIDs is compared to their adverse effect profiles.

### 4.2 Clinical Experience of Toxicity with Established NSAIDs

The most troublesome adverse effect associated with NSAID usage is gastrointestinal toxicity and it is this adverse effect that will be discussed in greatest detail.

Langman et al.<sup>[32]</sup> compared the use of NSAIDs in 1144 patients aged  $\geq 60$  years admitted to hospital with bleeding peptic ulcer with age- and gender-matched hospital and community controls. They

Table 1. Comparison of the cyclo-oxygenase (COX) selectivity of NSAIDs with the reported incidence (odds ratio) of upper gastrointestinal bleeding derived from 2 large case controlled studies<sup>[32,33]</sup> and their toxicity index<sup>[34]</sup>

NSAID	Odds ratio <sup>[32]</sup>	Odds ratio <sup>[33]</sup>	Toxicity index <sup>[34]</sup>	COX-2:COX-1 <sup>[47]</sup>	COX-2:COX-1 <sup>[34]</sup>	COX-2:COX-1 <sup>[31]</sup>
Azapropazone	31.5 (10.3-96.9)	23.4 (6.9-79.5)				
Ketoprofen	23.7 (7.6-74.2)	5.4 (2.6-11.3)	4.69			4.6
Piroxicam	3.7 (1.1-26.3)	18 (8.2-39.6)	3.96	33	9.54	
Indomethacin	11.3 (6.3-20.3)	6.3 (3.3-12.2)	5.15	107	22.3	14.7
Naproxen	9.1 (5.5-15.1)	3.1 (1.7-5.9)	3.01			3.3
Diclofenac	4.2 (2.6-6.8)		4.48	2.2		1.6
Ibuprofen	2.0 (1.4-2.8)	2.9 (1.7-5)	2.68		0.67	0.6
Tenoxicam				15		
Meloxicam				0.33		
Aspirin (acetylsalicylic acid)			1.77			
6-Methoxy-2-naphthylacetic acid <sup>a</sup>					0.14	0.3

a 6-Methoxy-2-naphthylacetic acid is the active metabolite of nabumetone.

found that peptic ulcer bleeding was strongly associated with the use of NSAIDs in the 3 months prior to admission. For the most commonly used NSAIDs it was possible to calculate an odds ratio for bleeding (table 1). Those NSAIDs with the least favourable odds ratios were azapropazone and ketoprofen, an intermediate ratio was seen for indomethacin, naproxen and piroxicam and the most favourable ratio was seen for ibuprofen and diclofenac. The authors noted that the risk of gastrointestinal bleeding for most NSAIDs tended to be greatest in those who had only recently started treatment. These patients were more likely to be on the full recommended dose than those who had been taking an NSAID long term.

Using a different approach to determine the risk of upper gastrointestinal bleeding with NSAID therapy, Garcia Rodriguez and Jick<sup>[33]</sup> identified 1457 cases of upper gastrointestinal bleeding from the records of general practitioners and calculated the odds ratios for gastrointestinal bleeding using appropriate controls (table 1). In general terms, the results give similar odds ratios for all gastrointestinal adverse effects.

A further measure of NSAID toxicity can be obtained from the ARAMIS (Arthritis, Rheumatism and Aging Medical Information System) database. The ARAMIS database contains detailed clinical information on more than 23 000 patients with rheumatic disorders in the US and Canada. The ARAMIS Post-Marketing Surveillance Program (PMS) has prospectively followed outcome status and drug adverse effects in a cohort of 2976 consecutively enrolled patients with rheumatoid arthritis giving 27 936 patient years of observation. Using the information obtained from this database Singh et al.<sup>[34]</sup> have developed a 'toxicity index' to give an overall measure of the adverse effect profile of different NSAIDs (and disease modifying agents which are not considered here).<sup>[34,35]</sup> The components of the index include adverse clinical symptoms, laboratory abnormalities and hospital admissions. Each adverse effect is weighted for severity based on the physician judgement. For each drug considered, the higher the toxicity index

the greater the drug toxicity (table 1). These 3 studies have provided an enormous amount of information about relative NSAID toxicity and it is of great interest to relate this information to their known COX selectivities.

The results of Langman et al.<sup>[32]</sup> and Garcia Rodriguez and Jick<sup>[33]</sup> are in reasonable agreement; both studies demonstrate that azapropazone, piroxicam and indomethacin have the highest odds ratios for gastrointestinal bleeding. The only major difference in the results of these 2 studies was for ketoprofen, which had a far less favourable odds ratio in the study of Langman et al.<sup>[32]</sup> compared with the odds ratio obtained by Garcia Rodriguez and Jick.<sup>[33]</sup>

Direct comparison between the odds ratios found in the studies of Langman et al.<sup>[32]</sup> and Garcia Rodriguez and Jick<sup>[33]</sup> with the toxicity index scores of Singh et al.<sup>[34]</sup> is difficult because this latter measure takes into account all adverse effects, not simply the incidence of gastrointestinal bleeding. Having said this, it can be seen that indomethacin and ketoprofen (according to the results of Langman et al.<sup>[32]</sup>) have both a high toxicity index and odds ratio for gastrointestinal bleeding. By contrast ibuprofen has a low toxicity index and odds ratio. A further point to note is the lower toxicity index associated with aspirin (acetylsalicylic acid) use than might have been expected when compared to the odds ratio.

The results of *in vitro* experiments for a particular NSAID cannot be directly translated into the clinical situation because of the differences in pharmacokinetic parameters to which the NSAID would be subjected. However, comparison of COX selectivities with adverse effects raises a number of points of interest.

- The COX selectivity of an individual NSAID is dependent on the biological assay used for its measurement. This means that the results of studies using different assays are not identical, but nevertheless the results do show a reasonable degree of agreement.
- The COX-1 selective drugs indomethacin and piroxicam are associated with an increased in-

cidence of gastrointestinal bleeding and high toxicity indices.

- Ibuprofen is more selective for COX-2 than COX-1 and is associated with a low incidence of gastrointestinal bleeding and low toxicity index.
- 6-Methoxy-2-naphthylacetic acid (6-MNA, the active metabolite of nabumetone) is more COX-2 than COX-1 selective. This NSAID is associated with a favourable adverse effect profile and this discussed further in section 5.1.
- Aspirin is the most COX-1 selective of the drugs tested and has a low toxicity index. This indicates that other factors are important in determining toxicity.
- Meloxicam is predominantly COX-2 selective. This new drug is discussed further in section 5.3.
- The COX selectivity of NSAIDs may be an indicator of the likelihood of adverse effects but the absolute concentration of NSAID required to inhibit COX is likely to be important also.

*In summary* the results of these clinical studies demonstrate convincingly that the incidence of gastrointestinal adverse effects varies widely between the currently available NSAIDs. COX-2 selectivity appears to be associated with a decreased incidence of gastrointestinal adverse effects.

## 5. New Developments In NSAID Therapy

There have been a number of new NSAIDs recently introduced onto the market which demonstrate favourable adverse effect profiles. It is interesting to investigate to what extent COX selectivity accounts for these properties.

### 5.1 Nabumetone

Nabumetone is a nonacidic NSAID that is established for the treatment of rheumatoid arthritis and osteoarthritis. Nabumetone can be distinguished from other NSAIDs by its excellent adverse effect profile.<sup>[36]</sup> Nabumetone has a number of interesting pharmacological properties which

collectively result in a reduced incidence of adverse effects.<sup>[37]</sup>

*In vitro* experimental work (see table 1) demonstrates that the active metabolite of nabumetone, 6-MNA, is COX-2 selective and this is likely to contribute to its favourable adverse effect profile.<sup>[31]</sup> Indeed using a human COX enzyme system Barnett et al.<sup>[31]</sup> found nabumetone to be approximately 3.5 times more active against COX-2 than COX-1. Nabumetone is nonacidic and not subject to anion trapping in the gastric mucosa, and it is a prodrug and as such the active drug does not come into direct with the gastric mucosa.

Nabumetone has been evaluated in clinical trials involving more than 7400 patients and post market surveillance data exists for more than 37 000 patients.<sup>[38]</sup> The cumulative incidence of nabumetone-induced gastric perforations, ulcers and haemorrhage lies between 0.3 and 0.95% and overall the withdrawal rate due to adverse effects lies between 3 and 13%.<sup>[38]</sup> Initial clinical studies involved more than 7000 patients in a number of double blind comparative drug trials (comparator drugs used were placebo, naproxen and aspirin), and during these trials, only 10 patients were admitted to hospital for reasons thought to be linked to nabumetone therapy.<sup>[39-41]</sup> Of these 10 patients, 6 had gastritis, 1 had abdominal pain, 1 had a gastric ulcer, 1 had biliary colic and 1 had a diffuse rash due to candidiasis.

In premarketing studies in the US, 930 patients were involved in double blind comparative trials (comparator drugs were placebo, naproxen and aspirin). Of these only 7 patients had evidence of gastrointestinal bleeding: 4 had a positive result to a guaiac test; 1 had a gastric ulcer; and 2 had bloody or black stools.<sup>[39-41]</sup>

When results from all the premarketing studies performed in the US are combined, data on nearly 2000 patients are available. From this data the cumulative incidence of peptic ulceration has been estimated to be 0.3% at 6 months (5 patients), 0.5% at 1 year (8 patients), 0.8% at 2 years (13 patients) and 0.95% at 8 years (16 patients). The average treatment duration before peptic ulcer development

was 24 months, a result that is not substantially different from the general population.

In the comparative short term studies, 1.4% of naproxen-treated patients developed an ulcer at 6 months.<sup>[38]</sup> The average time to ulcer development was 2.5 months. In these studies, the withdrawal rate during the first year of treatment was 13%. In a subsequent 8 year open label follow up evaluation the withdrawal rate was 3 to 6% for each subsequent year.<sup>[42]</sup>

In premarketing studies performed outside the US the overall withdrawal rate due to adverse events was 6%.<sup>[38]</sup> The most common adverse events were gastrointestinal (nausea, dyspepsia, diarrhoea, constipation and abdominal pain), skin rashes, headaches and dizziness). Bone marrow suppression, renal failure and severe dermatological or CNS conditions were not seen.

In a prospective postmarketing surveillance study performed in the UK, primary care physicians enrolled 10 800 patients who were being treated with nabumetone.<sup>[43]</sup> In addition, hospital physicians enrolled 2005 patients who had received other NSAIDs (ibuprofen, naproxen, diclofenac, piroxicam, ketoprofen, indomethacin or fenbufen) who acted as controls. NSAID therapy was being given to treat a number of musculoskeletal conditions. The patients in the nabumetone-treated group were followed up for 12 months and patients in the control group were only followed for 3 months.<sup>[43]</sup> During the study period there were 2 deaths in the nabumetone-treated group which may have been attributable to this therapy. One death was due to a gastric ulcer and the other secondary to renal failure in a patient known to have preexisting renal failure. The cumulative incidence of gastric perforations, ulcers and bleeds was 0.2% at 12 months. Renal insufficiency occurred in 9 patients (0.08%); in 2 patients the renal failure was reversible after cessation of therapy. Overall, the incidence of adverse events was similar between nabumetone and the comparator drugs, though the different lengths of follow-up need to be borne in mind. The subsequent withdrawal rate was approximately 10% in the nabu-

metone patients. The incidence of adverse effects increased when the nabumetone dosage was increased from 500 mg/day to 1 g/day daily. However, there was no further increase in adverse effect incidence when the dosage was increased further to 2 g/day.

In a postmarketing surveillance study conducted in Germany, follow-up data was obtained on 28 000 patients treated with nabumetone for 6 weeks.<sup>[37]</sup> During this short period there were no deaths attributable to nabumetone. Again the most common adverse effects were gastrointestinal. In total, 5 patients required hospital admission for gastrointestinal adverse effects (1 patient with a gastric ulcer, 1 with a duodenal ulcer, 2 with bleeding ulcers and 1 with an intestinal bleed).

In summary, the incidence of gastric perforations, ulcers or bleeds observed in nabumetone-treated patients who have taken part in premarketing and postmarketing studies has varied between 0.02 to 0.95%. This contrasts with the incidence of serious gastropathy associated with chronic usage of other NSAIDs of 2 to 4% per year.<sup>[44,45]</sup> The incidence of renal toxicity is also slight and nabumetone has demonstrated an adverse effect on renal function in less than 1% of patients.<sup>[46]</sup> These data thus demonstrate that nabumetone has a distinct tolerability advantage over other currently available NSAIDs.

## 5.2 Etodolac

The NSAID etodolac is established in the symptomatic treatment of rheumatoid arthritis and osteoarthritis (in the US it is licensed for the treatment of osteoarthritis alone). In clinical trials, etodolac has an efficacy equivalent to other NSAIDs, though the withdrawal rate due to gastrointestinal adverse effects has been less at 7%, which is considerably lower than the rates of 12 to 15% observed for aspirin and indomethacin.<sup>[47]</sup> In a study of patients with rheumatoid arthritis, treatment for 4 weeks with etodolac did not produce any significant change in gastric and duodenal prostaglandin levels obtained from biopsy specimens; significant changes were seen with the comparative drug

naproxen.<sup>[48]</sup> In animal studies etodolac has not been shown to deplete renal PGE<sub>2</sub>, in contrast to aspirin, indomethacin and piroxicam.<sup>[49]</sup> These results suggest COX-2 selectivity, a finding supported by the results of *in vitro* experimentation.<sup>[50]</sup>

### 5.3 Meloxicam

Meloxicam is a new NSAID of the enolic class which includes piroxicam and tenoxicam. It has recently been launched for the treatment of rheumatoid arthritis and osteoarthritis in South Africa and is approved in France, Sweden and Switzerland. In table 1 it can be seen that meloxicam is a more potent inhibitor of COX-2 than COX-1.<sup>[28]</sup>

Meloxicam is given once daily at a dose of 7.5 to 15 mg. At this dose the drug has linear pharmacokinetics and an elimination half-life of approximately 20 hours.<sup>[51]</sup> This short half-life (in contrast to piroxicam and tenoxicam with elimination half-lives of 50 and 70 hours respectively) reduces the chance of drug accumulation and may translate into an improved adverse effect profile. The drug has a bioavailability of 89% by the enteral route and effectively penetrates the inflamed rheumatoid synovium to achieve a concentration of approximately 50% of that seen in the plasma.<sup>[52,53]</sup> Animal studies have demonstrated almost complete metabolism with 30% biliary and 70% renal excretion. Hepatic metabolism is achieved, at least in part, by the cytochrome P450 system and it can be increased by enzyme inducers.<sup>[51]</sup>

In biological assays meloxicam has demonstrated the greatest COX-2 selectivity of the NSAIDs tested (table 1).<sup>[28]</sup> Indeed, in a study using a rat pleurisy model, meloxicam was twice as potent as tenoxicam, 3 times as potent as flubiprofen, 8 times as potent as diclofenac, and 20 times as potent as tenidap at inhibition PGE<sub>2</sub> biosynthesis.<sup>[54]</sup>

Meloxicam has now been administered to more than 5000 patients predominantly with rheumatoid arthritis or osteoarthritis in clinical trials.<sup>[55]</sup> The most commonly used comparator drugs used in these trials were diclofenac, piroxicam and naproxen. At a dosage of up to 15 mg/day meloxicam

has an efficacy equivalent to the comparative drugs with a trend in favour of improved gastrointestinal tolerability.<sup>[55]</sup> In the earliest stages of the drugs development doses up to 60 mg/day were used and these doses were not associated with increased gastrointestinal adverse effects.<sup>[55]</sup> This suggests that the conventional dose for meloxicam has a reasonable safety margin.

In the treatment of osteoarthritis meloxicam has been used in 5 major double blind trials involving about 1800 patients.<sup>[55-57]</sup> In these trials meloxicam demonstrated a clear superiority over placebo and an equivalent efficacy to comparator therapies. There was no significant difference in gastrointestinal adverse effects between therapies though the trend was in favour of meloxicam.

In the treatment of rheumatoid arthritis, controlled double blind studies over periods of 3 weeks to 6 months have demonstrated that the efficacy of meloxicam 7.5 mg/day and 15 mg/day is comparable to naproxen 750 mg/day<sup>[58]</sup> and piroxicam 20 mg/day, respectively.<sup>[59]</sup> In the 6 month comparison of meloxicam 7.5 mg/day (199 patients) with naproxen 750 mg/day (180)<sup>[58]</sup> there was no difference between the groups regarding the primary efficacy variables (global efficacy assessment by the patient and investigator, number of painful/tender and swollen joints). There were fewer gastrointestinal adverse effects in the meloxicam treated group (30%) compared with the naproxen-treated group (44%) and significantly more patients discontinued their therapy in the naproxen-treated group due to these adverse effects. In addition there were significant decreases in haemoglobin level and increases in serum creatinine and urea levels in the naproxen-treated group compared with the meloxicam group. However, the number of discontinuations due to lack of efficacy favoured the naproxen-treated group (26 patients) compared with the meloxicam-treated group (47).

In the 3-week comparison of meloxicam 15 mg/day (141 patients) with piroxicam 20 mg/day (135) there were no differences between the groups in terms of both efficacy and safety.<sup>[59]</sup> There was

a trend in favour of meloxicam with respect to gastrointestinal tolerance. There was no difference between the groups in terms of withdrawals due to lack of efficacy.

Overall, using data from all the above studies, the incidence of serious gastrointestinal events (perforation, ulcer and haemorrhage) was significantly lower in those patients taking meloxicam than diclofenac or piroxicam. For other body systems, meloxicam would appear to have a comparable safety profile to standard NSAIDs.<sup>[58-60]</sup>

*In summary*, meloxicam is a new NSAID which *in vitro* demonstrates high COX-2 selectivity. In the clinical trials thus far performed meloxicam appears to have an equivalent efficacy to standard NSAIDs combined with a lesser incidence of gastrointestinal adverse effects. The mechanism for this is not known but may relate to its COX-2 selectivity.

## 6. Conclusions

From the clinical data currently available it does seem that the COX selectivity of a NSAID is a significant determinant of the incidence of adverse effects. However, COX selectivity is not the whole story and other factors are also important. Currently, COX selectivity does not seem to be a clear determinant of clinical efficacy. It can be argued that of the currently available NSAIDs, nabumetone and etodolac have among the most favourable adverse effect profiles. Both of these drugs appear relatively COX-2 selective but have other important properties that contribute to their overall adverse effect profile. In initial studies, the new COX-2 selective inhibitor meloxicam, recently launched in its first market, has demonstrated clinical efficacy and a favourable gastrointestinal adverse effect profile.

Caution must remain about the potential of highly selective COX-2 inhibitors to affect normal physiological processes other than inflammation. It is interesting to note that mice genetically targeted to be deficient in COX-2 developed renal dysplasia (100% penetrance) and cardiac fibrosis (50% penetrance) but had normal inflammatory

responses to standard models.<sup>[61]</sup> The conclusion to be drawn is that the complete implications of COX inhibition are not yet understood. However, meloxicam, the most selective COX-2 NSAID discussed, has not been associated with an excess of any renal toxicity.

COX-2 inhibition may interfere with the normal physiology of the uterus and embryonic membranes, although clinically this should not pose a problem as NSAIDs are normally avoided during pregnancy. The clinical significance of inducible COX-2 in prostaglandin signalling in the brain is not known, hence the significance of inhibition is not known either. In the vascular endothelium and skeletal muscle the inducible enzyme appears to have an important role in the response to injury or mechanical stress. Again the significance of inhibition of these functions is not clear. The full effects of COX-2 inhibition can probably only be determined with accurate postmarketing surveillance studies.

## References

1. Vane JR. Inhibition of prostaglandin synthesis as a mechanism of action of aspirin like drugs. *Nature* 1971; 231: 235-7
2. Smith WL, Borgeat P, Fitzpatrick FA. The eicosanoids: cyclooxygenase, lipoxygenase, and epoxigenase pathways. In: Vance DE, Vance J, editors. *Biochemistry of lipids, lipoproteins and membranes*. Oxford: Elsevier Science Publishers, 1991: 297-325
3. Solomon LM, Juhlin L, Kirchenbain MB. Prostaglandins on cutaneous vasculature. *J Invest Dermatol* 1968; 51: 280-2
4. Ferreira SH. Prostaglandins, aspirin like drugs and analgesia. *Nature* 1992; 240: 200-3
5. Appleton I, Tomlinson A, Mitchell JA, et al. Distribution of cyclooxygenase isoforms in murine chronic granulomatous inflammation. Implications for future anti-inflammatory therapy. *J Pathol* 1995; 176 (4): 413-20
6. Dunn MJ. Non steroidal anti inflammatory drugs and renal function. *Annu Rev Med* 1984; 35: 411-28
7. Miller TA. Protective effects of prostaglandins against mucosal damage: current knowledge and proposed mechanisms. *Am J Physiol* 1983; 245: G601-23
8. Bunting S, Moneada S, Vane JR. The prostacyclin-thromboxane balance: pathophysiological and therapeutic considerations. *Br Med Bull* 1983; 39: 271-6
9. O'Banion MK, Sadowski HB, Winn V, et al. A serum and glucocorticoid regulated 4-kilobase mRNA encodes a cyclooxygenase related protein. *J Biol Chem* 1991; 266: 23261-7
10. Kujubu DA, Herschman HR. Dexamethasone inhibits nitrogen induction of the TIS10 prostaglandin synthase/cyclo-oxygenase gene. *J Biol Chem* 1992; 267: 7991-4
11. Sirois J, Richards JS. Purification and characterisation of a novel, distinct isoform of prostaglandin endoperoxide synthase induced by human chorionic gonadotrophin in

- granulosa cells of rat pre ovulatory follicles. *J Biol Chem* 1992; 267: 6382-8
12. Appleton I, Tomlinson A, Willoughby DA. Inducible cyclooxygenase (COX2): a safer therapeutic target? *Br J Rheumatol* 1995; 33 (5): 410-2
  13. Mead EA, Smith WL, De Witt DL. Differential inhibition of prostaglandin endoperoxide synthase (cyclooxygenase) isozymes by aspirin and other non steroidal anti inflammatory drugs. *J Biol Chem* 1992; 268: 6610-4
  14. Xie W, Chipman JG, Robertson DL, et al. Expression of a mitogen-responsive gene encoding prostaglandin synthase is regulated by mRNA splicing. *Proc Natl Acad Sci USA* 1991; 88: 2692-6
  15. Crofford LJ, Wilder RL, Ristimaki AP, et al. Cyclooxygenase 1 and 2 expression in rheumatoid synovial tissues: effect of interleukin 1 $\beta$ , Phorbol esters, and corticosteroids. *J Clin Invest* 1994; 93: 1095-101
  16. Knott I, Dieu M, Burton M, et al. Induction of cyclooxygenase by interleukin 1: comparative study between human synovial cells and chondrocytes. *J Rheumatol* 1994; 21: 462-6
  17. Schoen RT, Vender RJ. Mechanism of nonsteroidal anti-inflammatory drug induced gastric damage. *Am J Med* 1989; 86: 449-57
  18. Blanco A, Habib A, Levy-Toledano S, et al. Involvement of tyrosine kinases in the induction of cyclooxygenase-2 in human endothelial cells. *Biochem J* 1995; 312 (Pt 2): 419-23
  19. Inoue H, Yokoyama C, Hara S, et al. Transcriptional regulation of human prostaglandin-endoperoxide synthase-2 gene by lipopolysaccharide and phorbol ester in vascular endothelial cells. Involvement of both nuclear factor for interleukin-6 expression site and cAMP response element. *J Biol Chem* 1995; 270 (42): 249675-1
  20. Rimarachin JA, Jacobson JA, Szabo P, et al. Regulation of cyclooxygenase-2 expression in aortic smooth muscle cells. *Arterioscler Thromb* 1994; 14 (7): 1021-31
  21. Akai Y, Homma T, Burns KD, et al. Mechanical stretch/relaxation of cultured rat mesangial cells induces protooncogenes and cyclooxygenase. *Am J Physiol* 1994; 267: C482-90
  22. Martin M, Neumann D, Hoff T, et al. Interleukin-1 induced cyclooxygenase 2 expression is suppressed by cyclosporin A in rat mesangial cells. *Kidney Int* 1994; 45: 150-8
  23. Yamagata K, Andreasson KI, Kauffman WE, et al. Expression of a mitogen-inducible cyclooxygenase in brain neurons: regulation by synaptic activity and glucocorticoids. *Neuron* 1993; 11 (2): 371-86
  24. Adams, Cillaco-Morase Y, De Belleruche J. Cyclooxygenase-2 induction in cerebral cortex: an intracellular response to synaptic excitation. *J Neurochem* 1996; 66 (1): 6-13
  25. Vanderburgh HH, Shansky J, Solerssi R, et al. Mechanical stimulation of skeletal muscle increases prostaglandin F2 alpha production, cyclooxygenase activity, and cell growth by a pertussis toxin sensitive mechanism. *J Cell Physiol* 1995; 163 (2): 285-94
  26. Jacobs AL, Hwang D, Julian J, et al. Regulated expression of prostaglandin endoperoxide synthase-2 by uterine stroma. *Endocrinology* 1994; 135: 1807-15
  27. Slater DM, Berger LC, Newton, et al. Expression of cyclooxygenase types 1 and 2 in human fetal membranes at term. *Am J Obstet Gynaecol* 1992; 172 (1 Pt 1): 77-82
  28. Englehardt G. Meloxicam: a potent inhibitor of COX2 [abstract]. 9th International Conference on Prostaglandins and Related Compounds: 1994 Jun 6-10; Florence: 82
  29. Meade EA, Smith WL, De Witt DL. Differential inhibition of prostaglandin endoperoxide synthase (cyclooxygenase) isozymes by aspirin and other non-steroidal anti-inflammatory drugs. *J Biol Chem* 1993; 268 (9): 6610-4
  30. Mitchell JA, Akaravereenont P, Thiemermann C, et al. Selectivity of nonsteroidal antiinflammatory drugs as inhibitors of constitutive and inducible COX1. *Proc Natl Acad Sci USA* 1994; 90: 11163-697
  31. Barnett J, Chow J, Ives D, et al. Purification, characterisation and selective inhibition of human prostaglandin G/H synthase 1 and 2 expressed in the baculovirus system. *Biochim Biophys Acta* 1994; 1209 (1): 130-9
  32. Langman MJS, Weil J. Risks of bleeding peptic ulcer associated with individual non-steroidal anti-inflammatory drugs. *Lancet* 1994; 343: 1075-8
  33. Garcia Rodriguez LA, Jick H. Risk of upper gastrointestinal bleeding and perforation associated with individual non-steroidal anti-inflammatory drugs. *Lancet* 1994; 343: 769-72
  34. Singh G, Ramey DR, Morfeld D, et al. Comparative toxicity of non-steroidal anti-inflammatory agents. *Pharmacol Ther* 1994; 62: 175-91
  35. Fries JF, Spitz PW, Williams CA, et al. A toxicity index for comparison of side effects among different drugs. *Arthritis Rheum* 1990; 34: 121-30
  36. Richardson CE, Emery P. Innovative treatment approaches for rheumatoid arthritis. *Baillieres Clin Rheumatol* 1995; 9 (4): 731-58
  37. Blower PR. The unique pharmacological profile of nabumetone. *J Rheumatol* 1992; 19 Suppl. 36: 13-9
  38. Gerson GC. Worldwide safety experience with nabumetone. *J Rheumatol* 1992; 19 Suppl. 36: 48-56
  39. Wilkens RF. An overview of the long term safety experience of nabumetone. *Drugs* 1990; 40 Suppl. 5: 34-7
  40. Fleishman RM. Clinical efficacy and safety of nabumetone in rheumatoid arthritis and osteoarthritis. *J Rheumatol* 1992; 19 Suppl. 36: 32-40
  41. Jackson RE, Mitchell FN, Brindley DA. Safety evaluation of nabumetone in US clinical trials. *Am J Med* 1987; 83 Suppl. 4B: 115-20
  42. Jenner PM, Johnson ES. Review of experience with nabumetone in clinical trials outside of the United States. *Am J Med* 1987; 83 Suppl. 4B: 110-4
  43. Inman WHW, Wilton WV, Pearce GI, et al. Prescription event monitoring with nabumetone. *Pharm Med* 1990; 4: 309-17
  44. Hochberg MC. Association of non steroidal anti-inflammatory drugs with upper gastrointestinal disease: epidemiological and economic considerations. *J Rheumatol* 1992; 19 Suppl. 36: 63-7
  45. Soll AH. Pathogenesis of peptic ulcer and implications for therapy. *N Engl J Med* 1990; 322: 909-16
  46. Arnoff GR. Therapeutic implications associated with renal studies of nabumetone. *J Rheumatol* 1992; 19 Suppl. 36: 25-31
  47. Schattenkirchner M. An updated safety profile of etodolac in several thousand patients. *Eur J Rheumatol Inflamm* 1990; 10: 56-65
  48. Tana AS, McLaughlin S, Holland PJ, et al. Effect on gastric and duodenal mucosal prostaglandins of repeated intake of therapeutic doses of naproxen and etodolac in rheumatoid arthritis. *Ann Rheum Dis* 1990; 49: 354-8
  49. Lee DKH. Effect of etodolac on the prostaglandin concentration in the kidney of the normal rat. *Drug Dev Res* 1986; 9 (4): 305-11
  50. Glaser K, Sung ML, O'Neill K, et al. Etodolac selectively inhibits human prostaglandin G/H synthase 2 (PGHS-2) versus PGHS1. *Eur J Pharmacol* 1995; 281: 107-11



51. Turek D, Roth W, Busch U. A review of the pharmacokinetics of meloxicam. *Br J Rheumatol* 1996; 35 Suppl.: 13-6
52. Degner F, Heinzel G, Busch U. Transsynovial kinetics of meloxicam [abstract]. *Scand J Rheumatol* 1994; suppl. 98: 121
53. Busch U, Englehart G. Distribution of [<sup>14</sup>C] meloxicam in joints of rats with adjuvant arthritis. *Drugs Exp Clin Res* 1990; 16: 49-52
54. Englehart JG, Bogel R, Schnitzler C, et al. Meloxicam: influence on arachidonic acid metabolism. Part II. In vitro findings. *Biochem Pharmacol* 1996; 51 (1): 29-38
55. Distel M, Mueller C, Bluhmki E, et al. Safety of meloxicam: a global analysis of clinical trials. *Br J Rheumatol* 1996; 35 Suppl. 1: 68-77
56. Linden B, Distel M, Bluhmki E. A double blind study to compare the efficacy and safety of meloxicam 15mg with piroxicam 20mg in patients with osteoarthritis of the hip. *Br J Rheumatol* 1996; 35 Suppl. 1: 35-8
57. Hosie J, Distel M, Bluhmki E. Meloxicam in osteoarthritis: a 6 month double blind comparison with diclofenac sodium. *Br J Rheumatol* 1996; 35 Suppl. 1: 39-43
58. Wojtulewski JA, Schattenkirchner M, Barcelo P, et al. A six month double blind trial to compare the efficacy and safety of meloxicam 7.5 mg daily and naproxen 750 mg daily patients with rheumatoid arthritis. *Br J Rheumatol* 1996; 35 Suppl. 1: 22-9
59. Huskisson EC, Narjes H, Bluhmki E. Efficacy and tolerance of meloxicam, a new NSAID in daily oral doses of 15, 30 and 60mg in comparison with 20mg piroxicam in patients with rheumatoid arthritis [abstract]. *Scand J Rheumatol* 1994; Suppl. 98: 115
60. Huskisson EC, Ghosian R, Kurthen R, et al. A long term study to evaluate the safety and efficacy of meloxicam therapy in patients with rheumatoid arthritis. *Br J Rheumatol* 1996; 35 Suppl. 1: 29-34
61. Dinchev JE, Car BD, Focht RJ, et al. Renal abnormalities and an altered inflammatory response in mice lacking cyclooxygenase II. *Nature* 1995; 378: 406-9

---

Correspondence and reprints: Dr Charles Richardson, Department of Rheumatology and Rehabilitation, University of Leeds, 36 Clarendon Road, Leeds LS2 9NZ, England.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**